

You Can Manage Only What You Measure!

Improving Treatment Effectiveness

- **Scours:** How long after birth are calves receiving their first feeding of colostrum?
- **Scours:** How much colostrum are calves receiving at their first feeding?

- **Scours:** Is the first-feeding colostrum uniformly of high quality?
- **Scours:** Is the first-feeding colostrum uniformly low in bacteria content as fed?

- **Scours:** Do we have a procedure (protocol) that everyone uses to diagnose treatable cases of scours (diarrhea)?
- **Scours:** Do we record calf ID, severity of symptoms, drug, dose and duration of treatment for each calf receiving antibiotics for diarrhea symptoms?

- **Scours:** Do we regularly summarize treatment records to identify the most effective treatment procedures?

- **Pneumonia:** Do we have a procedure that everyone uses to diagnose respiratory illness (pneumonia)?
- **Pneumonia:** Do we record calf ID, severity of symptoms, drug, dose and duration of treatment for each calf treated for respiratory illness?

- **Pneumonia:** Do we regularly summarize treatment records to identify the most effective treatment procedures?

Do you think that the title of this issue is probably an overstatement? I am going to argue that when you do not have the facts about your calf enterprise you are only “muddling through” rather than managing.

Example Number 1: Managing scours (diarrhea)

How much do we really know about our calf enterprise? How many calves are getting their first feeding of colostrum within the first few hours of life (percent)? Is this written down anywhere? If we keep this record, how recently have we summarized these facts so that they are usable information? Remember, we lose about forty percent of the calf’s ability to absorb antibodies when the first feeding comes later than six hours after birth.

Sam Leadley, Calf & Heifer Management Specialist

sleadley@yahoo.com www.atticacows.com

For Calves with Sam blog go to dairycalfcare.blogspot.com

© Attica Vet. Assoc. 2017 All Rights Reserved.

How much colostrum are the calves actually consuming? For example, the farm protocol may say that every calf receives four quarts of colostrum in the first six hours. How closely is this procedure being followed? In practice, are the calves born at night assumed to have nursed? And, then these calves are not fed any colostrum until the next regular calf feeding time? Remember, calves cannot absorb antibodies that have not been fed.

Are we assuming that all colostrum that “looks good” has an adequate concentration of antibodies for successful passive transfer of immunity? Even in a well-vaccinated herd it has been estimated that as many as one-fourth of the cows give poor quality colostrum. While the low quality colostrum is great nutritionally, it does need to be supplemented if that is all available to be fed. Remember, for a given volume fed, antibody transfer works better with high rather than low quality colostrum. Quality can be checked on-farm very quickly using a Brix refractometer - [Read More Here](#).

Does our calf enterprise track colostrum cleanliness? This is a key piece of information for quality control in calf care. Recent experiences in a number of states emphasize for me the universal challenge of feeding clean colostrum. Even samples from well-managed dairy farms will be contaminated with enough coliform bacteria to predict treatable cases of scours. As an aside, this is a challenge when feeding colostrum replacer, too.

If your farm has not submitted colostrum samples for bacterial culturing within the past six months, now is the time to do it. Using a sterile milk sample bottle (just like the ones used by the milk truck driver), from the end of a tube feeder or nipple bottle just as you are ready to feed colostrum to a newborn calf collect one-half a sample vial of colostrum. Freeze it. Have it cultured asking for both speciation and quantification. Coliform bacteria counts over 5,000cfu/ml are not good – for a resource on bacteria standards click [HERE](#).

Remember, contaminated colostrum hits calves twice. First, they may transfer some the bacteria directly into their blood. Second, many bad bacteria will attach to the lining of the gut ready to make trouble for weeks.

To achieve uniformity in scours diagnosis it may be helpful to use the chart at this web site: http://www.vetmed.wisc.edu/dms/fapm/fapmtools/8calf/calf_health_scoring_chart.pdf . Scroll to the bottom of the second page for scours pictures.

Improving the effectiveness of our treatment protocols is a great goal. This not easy however.

Example Number 2: Calves treated for pneumonia.

Are too many calves being treated for pneumonia? How effective is the treatment protocol? How many calves treated for pneumonia are requiring re-treatment?

Do we have a procedure that everyone uses when diagnosing pneumonia? If you have not given this much thought, consider Dr. McGuirk’s (University of Wisconsin) diagnostic scoring

Sam Leadley, Calf & Heifer Management Specialist

sleadley@yahoo.com www.atticacows.com

For Calves with Sam blog go to dairycalfcare.blogspot.com

© Attica Vet. Assoc. 2017 All Rights Reserved.

system with pictures. It is located at this web site: http://www.vetmed.wisc.edu/dms/fapm/fapmtools/8calf/calf_respiratory_scoring_chart.pdf See the second page for pictures of healthy and sick calves. Early diagnosis of pneumonia and timely treatment are keys to full recovery. Also, it is a best management practice to work with the herd veterinarian to set farm-specific thresholds for starting treatment of sick calves.

How complete are our records? We need to write down details for each calf diagnosed with pneumonia [for example, ID, severity of symptoms, treatment given]. These facts along with the age of the calf when she was treated need to be summarized regularly to give us useful information. Examples are:

- Percent calves treated one or more times between birth and weaning.
- Percent calves being retreated for pneumonia.
- For an individual antibiotic, percent calves successfully recovering from pneumonia.

I like to summarize treatments by age, too.

- In the winter many cases around two weeks of age may point to inadequate energy from milk or milk replacer for environmental conditions. Too little energy for too long results in low resistance to pneumonia.
- Many cases around four weeks old may point to inadequate coccidia control. Subclinical coccidiosis leads to low resistance to pneumonia.
- Many pneumonia treatments right around weaning suggests inappropriate weaning management.

Bottom line?

Collecting and recording good reliable facts and summarizing them to make useful information leads to good management. We manage only what we measure.